

Buckeye Butterfly

Class	Order	Family	Species
Insecta	Lepidoptera	Nymphalidae	Junonia coenia

Range

Growth: Egg:

complete metamorphosis

Caterpillar: 2-4 broods per year. 3-4 weeks

Pupa: 2 weeks

Adult:

overwinters as an adult

Habitat

Most of North

America and

worldwide.

Open, neglected fields, grasslands, temperate and tropical zones.

Niche

Diurnal, spring to fall breeder, weedy roadsides and fields.

Diet

Nature: caterpillar-plantain (Plantago spp.), snapdragon, monkey flower (Mimulus) adult - flower nectar Captivity: caterpillar - same as above adult - sugar water

Reproduction

laid from early spring to early fall; one week

Physical Characteristics

Adult Larvae

Mouthparts: siphoning/sucking chewing 3 pair, 5 pair fleshy prolegs 6 legs Legs:

Wings: none 2 pair

roundish, ribbed and cryptically colored green Eggs: Larvae: black, very spiny

Pupae: (chrysalis): mottled pale brown Adult: brown marked by a large and small eye spot and

orange bars near the wing edge. 2.5 inch wing

spread.

Social Organization

None

Special Adaptations

Morphology: The Nymphalid family of butterflies are all colored in brown, orange and black.

Sources

(5)(6)(27)

